



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A23J 3/34, A23K 1/14, A23L 1/105	A1	(11) International Publication Number: WO 95/28850 (43) International Publication Date: 2 November 1995 (02.11.95)
(21) International Application Number: PCT/DK95/00166 (22) International Filing Date: 20 April 1995 (20.04.95) (30) Priority Data: 470/94 22 April 1994 (22.04.94) DK (71) Applicant (for all designated States except US): NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): NIELSEN, Per, Munk [DK/DK]; Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd (DK). KNAP, Inge, Helmer [DK/DK]; Novo Nordisk A/S, Novo Allé, DK-2880 Bagsværd (DK). (74) Common Representative: NOVO NORDISK A/S; Corporate Patents, Novo Allé, DK-2880 Bagsværd (DK).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>With amended claims.</i>	
(54) Title: A METHOD FOR IMPROVING THE SOLUBILITY OF VEGETABLE PROTEINS (57) Abstract <p>The present invention relates to a method for improving the solubility of vegetable proteins. More specifically, the invention relates to methods for the solubilization of proteins in vegetable protein sources, which methods comprise treating the vegetable protein source with an efficient amount of one or more phytase enzymes, and treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes. In another aspect, the invention provides animal feed additives comprising a phytase and one or more proteolytic enzymes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

A METHOD FOR IMPROVING THE SOLUBILITY OF VEGETABLE PROTEINS

TECHNICAL FIELD

The present invention relates to a method for improving the solubility of vegetable proteins. More specifically, the invention relates to methods
5 for the solubilization of proteins in vegetable protein sources, which methods comprise treating the vegetable protein source with an efficient amount of one or more phytase enzymes, and treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes. In another aspect the invention provides animal feed additives comprising a phytase and one or more proteolytic
10 enzymes.

BACKGROUND ART

Protein is an essential nutritional factor for a.o. mammals and layers. Most livestock and many human beings get the necessary proteins from vegetable protein sources. Important vegetable protein sources are e.g. cereals,
15 legumes and oilseed crops.

Hydrolyzed vegetable proteins, such as soy protein hydrolysates, find application as nutrients, e.g. as nutritional additives to foods and beverages. Hydrolyzed proteins are absorbed more easily than unhydrolysed protein, why protein hydrolysates are considered having the highest nutritional value.
20 Several methods for preparing protein hydrolysates are known and described in the literature, cf. e.g. US 4,324,805 and WO 92/15696.

Essentially all food and feed substances originating from plants contain phytate and phytic acid as a storage phosphorus source [cf. review in *E. Graf (ed.), Phytic Acid, Chemistry and Applications*, Minneapolis, U.S.A., 1986].
25 About 75-78% of the phosphorus in cereals is bound as phytic acid. Phytate comprises 1-3% of all nuts, cereals, legumes, oil seeds, spores and pollen. Complex salts of phytic acid are termed phytin.

Phytic acid chelates minerals such as calcium, zinc, magnesium, and iron, thereby decreasing the bio-availability of nutritionally important minerals, and is generally considered an anti-nutritional factor. In vitro studies indicate that phytic acid inhibits the peptic digestion of some animal proteins, whereas the trypsin digestion was unaffected [cf. *Knuckles et al.*, Journal of Food Science, 1989 54 1348-1350].

Phytases are enzymes which catalyze the conversion of phytate to inositol and inorganic phosphorus. Phytases have been obtained from e.g. *Bacillus*, *Pseudomonas*, *Saccharomyces* and *Aspergillus*.

10 In US 3,297,548 it has been suggested to add microbial phytase to feedstuffs of monogastric animals in order to avoid supplementing the feed with inorganic phosphorus.

It has also been conceived that phytases may be used in the processing of soy. Thus EP-A-0 420 358 reports that soybean meal contains high 15 levels of the anti-nutritional factor phytate which renders this protein source unsuitable for application in baby food, as well as in feed for fish, calves and other non-ruminants, since the phytate chelates essential minerals present therein.

In summary it has previously been suggested to use phytase enzymes either for exploiting the phosphorus bound in the phytate/phytic acid 20 present in vegetable protein sources, or for exploiting the nutritionally important minerals bound in phytic acid complexes.

However, there is a need for improving the availability of the proteins present in vegetable sources, e.g. cereals, legumes and oilseed crops, since an increased availability leads to higher yields of protein hydrolyzing processes 25 as well as nutritional benefits, i.e. improved protein utilization.

SUMMARY OF THE INVENTION

It has now been found that the solubility of proteins present in vegetable sources may be increased by treating the vegetable source with an efficient amount of a phytase enzyme. By addition of a phytase during a proteolytic

process, higher degrees of hydrolysis and improved protein solubility are obtained, and yields are improved.

Accordingly, the present invention provides a method for solubilizing proteins in a vegetable protein source, which method comprises treating the vegetable protein source with an efficient amount of one or more phytase enzymes, and treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes.

In another aspect, the invention provides an animal feed additive comprising a phytase and a proteolytic enzyme.

10

DETAILED DISCLOSURE OF THE INVENTION

Solubilization of Vegetable Proteins

The present invention provides a method for the solubilization of proteins in a vegetable protein source, thereby obtaining a protein hydrolysate. The method comprises the steps of

15 (a) treating the vegetable protein source with an efficient amount of one or more phytase enzymes; and

(b) treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes.

Compared to known method for obtaining protein hydrolysates, 20 the present method improves the availability of the proteins, thereby leading to increased extraction, higher yields and improved protein utilization.

The vegetable protein source subjected to the method of the invention may be any proteinaceous vegetable, in particular a legume, a cereal, a composite plant or a crucifera. Preferably, the legume is selected from the group 25 consisting of soy bean, faba bean, pea, and lupine (*Lupinus albus*), more preferably the legume is soy beans. Preferably, the cereal is selected from the group consisting of wheat, corn, barley, rye, oat, rice, sorghum, sesame (*Sesamum indicum*), and millet (*Panicum miliaceum*). An example of a useful composite plant is sunflower

(*Helianthus*), and an example of a useful crucifera is rape (rape seed, e.g. *Brassica napus*).

The vegetable protein source is preferably soy bean.

The proteinaceous vegetable subjected to the method of the invention may be provided in any form, including "processed forms" wherein the protein content of the dry matter is increased before the protein subjected to the method according to the invention. For instance, soy protein raw material may obtained as soy beans, defatted soy flakes, soy meal, soy concentrate or soy isolate. Such proteinaceous raw material typically contains from about 42% to about 95% protein.

10 The protein source may be pre-treated in any conventional manner prior to being subjected to the method of the present invention. In particular the protein source may be dried and/or grounded, i.e. processed into meal or flakes (such as soy bean meal or flakes).

The two steps, (a) and (b) of the method of the invention, may 15 be carried out as consecutive steps, or the two steps may be carried out simultaneously.

Also, the vegetable protein source may be treated with one enzyme preparation comprising one or more phytase enzymes and one or more proteolytic enzymes, or the vegetable protein source may be treated with two or 20 more enzyme comprising a one or more phytase enzymes and/or one or more proteolytic enzymes.

It is at present contemplated that the method of the invention is preferably carried out at a pH of the suspension of from about 3 to about 9, more preferably of from about pH 4 to about pH 7.

25 It is also contemplated that the method of the invention is preferably carried out at a temperature of from about 10°C to about 70°C, more preferably of from about 35°C to about 65°C.

In a more specific embodiment, the method comprises the subsequent steps of

30 (i) suspending a vegetable protein source in water;

(ii) proteolytically hydrolysing the protein by subjecting the suspension to enzymatic treatment with one or more phytase enzymes and one or more proteolytic enzyme;

(iii) inactivating the enzymes; and optionally

5 (iv) separating the hydrolysed protein from the suspension.

Inactivation of the enzyme may be carried out by conventional methods for inactivating enzymes, e.g. by heat treatment, i.e. elevating the temperature of the hydrolysis suspension or mixture to a temperature which denatures the enzymes, typically to a temperature of above 85°C.

10 The separating step may be carried out by conventional methods for separating hydrolysed protein from suspensions, e.g. by centrifugation or membrane filtrating the suspension containing the protein hydrolysate.

In general terms, the method of the invention may be carried out similar to conventional methods for obtaining protein hydrolysates, e.g. as
15 described in US 4,324,805, US 4,100,024 and WO 92/15696, which publications are hereby incorporated by reference.

Phytase Enzymes

The method of the invention for the solubilization of proteins in a vegetable protein source comprises treating the vegetable protein source with
20 an efficient amount of one or more phytase enzymes.

In the context of this invention, a phytase enzymes is an enzyme which catalyzes the removal of inorganic phosphorous from various myoinositol phosphates. Phytase enzymes are preferably derived from a microbial source such as bacteria, fungi and yeasts, but may also be of vegetable origin.

25 In a preferred embodiment, the phytase enzyme is derived from a fungal strain, in particular a strain of *Aspergillus*, e.g. *Aspergillus niger*, *Aspergillus oryzae*, *Aspergillus ficuum*, *Aspergillus awamori*, *Aspergillus nidulans* and *Aspergillus terreus*. Most preferred is a phytase enzyme derived from a strain of *Aspergillus niger* or a strain of *Aspergillus oryzae*.

In another preferred embodiment, the phytase enzyme is derived from a bacterial strain, in particular a strain of *Bacillus* or a strain of *Pseudomonas*. Preferably the phytase enzyme is derived from a strain of *Bacillus subtilis*.

In yet another preferred embodiment, the phytase enzyme is derived from a yeast, in particular a strain of *Kluveromyces* or a strain of *Saccharomyces*. Preferably the phytase enzyme is derived from a strain of *Saccharomyces cerevisiae*.

In the context of this invention "an enzyme derived from" encompasses an enzyme naturally produced by the particular strain, either recovered from that strain or encoded by a DNA sequence isolated from this strain and produced in a host organism transformed with said DNA sequence.

The phytase enzyme may be derived from the microorganism in question by use of any suitable technique. In particular, the phytase enzyme may be obtained by fermentation of a phytase producing microorganism in a suitable nutrient medium, followed by isolation of the enzyme by methods known in the art.

The broth or medium used for culturing may be any conventional medium suitable for growing the host cell in question, and may be composed according to the principles of the prior art. The medium preferably contain carbon and nitrogen sources and other inorganic salts. Suitable media, e.g. minimal or complex media, are available from commercial suppliers, or may be prepared according to published receipts, e.g. the American Type Culture Collection (ATCC) Catalogue of strains.

After cultivation, the phytase enzyme is recovered by conventional method for isolation and purification proteins from a culture broth. Well-known purification procedures include separating the cells from the medium by centrifugation or filtration, precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, and chromatographic methods such as e.g. ion exchange chromatography, gel filtration chromatography, affinity chromatography, etc.

Alternatively, the phytase enzyme is preferably produced in larger quantities using recombinant DNA techniques, e.g. as described in EP-A1-0 420 358, which publication is hereby incorporated by reference.

Preferably, a fungus of the species *Aspergillus* which has been transformed with the phytase-encoding gene obtained from the species *Aspergillus ficuum* or *Aspergillus niger*, is cultured under conditions conducive to the expression of the phytase-encoding gene as described in EP-A1-0 420 358.

The phytase-containing fermentation broth is preferably treated by means of both filtration and ultra-filtration prior to being used in the (formulation) of the present invention.

It is at present contemplated that an amount of phytase efficient for improving the solubility of the vegetable protein corresponds to of from about 2 to about 50000 FYT (as defined below) per kg of vegetable protein source, preferably of from about 50 to about 30000 FYT per kg of vegetable protein source, most preferred of from about 100 to about 10000 FYT per kg of vegetable protein source.

Proteolytic Enzymes

The method of the invention for the solubilization of proteins in a vegetable protein source also comprises treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes.

The proteolytic enzyme may be a microbial enzyme, preferably a protease derived from a bacterial or a fungal strain, or the proteolytic enzyme may be trypsin or pepsin. In a preferred embodiment, the proteolytic enzyme is a bacterial protease derived from a strain of *Bacillus*, preferably a strain of *Bacillus subtilis* or a strain of *Bacillus licheniformis*. Commercially available *Bacillus* proteases are Alcalase™ and Neutrase™ (Novo Nordisk A/S, Denmark). In another preferred embodiment, the proteolytic enzyme is a fungal protease derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus aculeatus*, a strain of *Aspergillus niger*, a strain of *Aspergillus oryzae*. A commercially available *Aspergillus* protease is Flavourzyme™ (Novo Nordisk A/S, Denmark).

It is at present contemplated that the amount of proteolytic enzyme efficient for improving the solubility of the vegetable protein corresponds to of from about 0.0001 to about 0.5 AU (as defined below) per kg of vegetable protein source, preferably of from about 0.001 to about 0.05 AU per kg of vegetable protein source, more preferred of from about 0.005 to about 0.03 AU per kg of vegetable protein source.

Additional Enzymes

As described above, the present invention provides a method for the solubilization of proteins in a vegetable protein source, which method
10 comprises the steps of

(a) treating the vegetable protein source with an efficient amount of one or more phytase enzymes; and

(b) treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes.

15 The two steps, (a) and (b) of the method of the invention, may be carried out as consecutive steps, or the two steps may be carried out simultaneously.

In a preferred embodiment, the method of the invention further comprises the step of

20 (c) treating the vegetable protein source with an efficient amount of one or more enzymes selected from the group consisting of lipolytic enzymes and glucosidase enzymes.

The three steps, (a), (b) and (c) of the method of the invention, may be carried out as consecutive steps, or the three steps may be carried out
25 simultaneously. Or the two steps, (a) and (b) of the method of the invention, may be carried out simultaneous, followed by step (c).

The lipolytic enzyme of step (c) may be any triacylglycerol lipase (EC 3.1.1.3).

The glycosidase enzyme of step (c) may be any glycosidase
30 enzyme (EC 3.2, also known as carbohydrases). Preferably, the glycosidase enzyme

is an amylase, in particular an α -amylase or a β -amylase, a cellulase, in particular an endo-1,4- β -glucanase (EC 3.2.1.4) or an endo-1,3- β -glucanase (3.2.1.6), a xylanase, in particular an endo-1,4- β -glucanase (EC 3.2.1.8) or a xylan-endo-1,3- β -xylosidase (EC 3.2.1.32), an α -galactosidase (EC 3.2.1.22), a polygalacturonase (EC 3.2.1.15, also known as pectinases), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91, also known as cellobiohydrolases), an endoglucanase, in particular an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

In a more preferred embodiment, the glucosidase enzyme is a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), or an endoglucanase, preferably an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

Industrial Applications

The method of the invention for the solubilization of proteins in a vegetable protein source, may find application in various industries. In particular the method of the invention find application for the preparation of protein hydrolysates useful as nutrients, e.g. as nutritional additives to foods and beverages.

Therefore, in another aspect, the present invention also provides a protein hydrolysate obtained from a vegetable protein source by the method of the invention.

The protein hydrolysate of the invention may be added to foods or beverages, thereby increasing the nutritional value.

In particular the protein hydrolysate of the invention may be added to animal feed, e.g. feedstuffs for monogastric animals.

Animal Feed Additives

The present invention provides an animal feed additive comprising one or more phytase enzymes and one or more proteolytic enzymes.

When added to animal feed, the combination of one or more phytase enzymes and one or more proteolytic enzymes improves the availability of the proteins present in animal feed of vegetable origin, thereby leading to increased extraction of the vegetable proteins, higher protein yields and improved protein utilization. Thereby, the nitrogen digestibility and the nutritive value of the fodder becomes increased, and the growth rate and/or feed conversion ratio (i.e. the weight of ingested feed relative to weight gain) of the animal is improved.

In the context of this invention, an animal feed additive is an enzyme preparation comprising one or more phytase enzymes and one or more proteolytic enzymes and suitable carriers and/or excipients, and which enzyme preparation is provided in a form that is suitable for being added to animal feed. The animal feed additive of the invention may be prepared in accordance with methods known in the art and may be in the form of a dry or a liquid preparation. The enzyme to be included in the preparation, may optionally be stabilized in accordance with methods known in the art.

In a specific embodiment the animal feed additive of the invention is a granulated enzyme product which may readily be mixed with feed components, or more preferably, form a component of a pre-mix. The granulated enzyme product may be coated or uncoated. The particle size of the enzyme granulates preferably is compatible with that of feed and pre-mix components. This provides a safe and convenient mean of incorporating the enzymes into feeds.

In another specific embodiment, the animal feed additive of the invention is a stabilized liquid composition, which may be an aqueous or oil-based slurry.

In yet another specific embodiment, one or more of the enzymes are applied before or after pelleting or extrusion of the feed.

The animal feed additive of the invention may exert its effect either *in vitro* (by modifying components of the feed) or *in vivo*. The feed additive of the invention is particularly suited for addition to animal feed compositions containing high amounts of proteinaceous vegetables, in particular legumes, cereals, composite plants or crucifera. Preferably, the legume is soy bean, faba bean, pea, and/or lupi-

ne (*Lupinus albus*). Most preferably the legume is soy beans. Preferably, the cereal is wheat, corn, barley, rye, oat, rice, sorghum, sesame (*Sesamum indicum*), and millet (*Panicum miliaceum*). An example of a composite plant is sunflower (*Helianthus*), and an example of a useful crucifera is rape (rape seed, e.g. *Brassica napus*).

5 In another preferred embodiment, the invention provides an animal feed additive which comprises one or more additional enzymes selected from the group consisting of lipolytic enzymes and glucosidase enzymes.

The lipolytic enzyme may be any triacylglycerol lipase (EC 3.1.1.3).

10 The glycosidase enzyme may be any glycosidase enzyme (EC 3.2, also known as carbohydrases). Preferably, the glycosidase enzyme is an amylase, in particular an α -amylase or a β -amylase, a cellulase, in particular an endo-1,4- β -glucanase (EC 3.2.1.4) or an endo-1,3- β -glucanase (3.2.1.6), a xylanase, in particular an endo-1,4- β -glucanase (EC 3.2.1.8) or a xylan-endo-1,3- β -xylosidase
15 (EC 3.2.1.32), an α -galactosidase (EC 3.2.1.22), a polygalacturonase (EC 3.2.1.15, also known as pectinases), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91, also known as cellobiohydrolases), an endoglucanase, in particular an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

20 In a more preferred embodiment, the glucosidase enzyme is a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), or an endoglucanase, preferably an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

25 It is at present contemplated that the pH of the animal feed additive should be in the range of from about pH 2 to about pH 8.

It is at present contemplated that the amount of phytase activity in the animal feed additive should be in the range of from about 200 to about 50000 FYT (as defined below) per gram of the total composition, preferably in the range
30 of from about 500 to about 10000 FYT per gram of the total composition, most

preferred in the range of from about 2000 to about 6000 FYT per gram of the total composition.

Preferably, the amount of additive added to the animal feed should be sufficient to provide a feed composition containing at least 50 FYT per kg feed, more preferably, between about 100 and about 2000 FYT per kg feed.

It is to be understood that the actual amount of Phytase Units which should be added to the feed in order to enhance the nutritive value, i.e. the nitrogen digestibility, will depend on the composition of the feed itself. Feedstuffs containing high amounts of phytic acid will generally require the addition of higher amounts of phytase activity. The necessary amount of phytase may easily be determined by the skilled person.

Phytase Activity (FYT)

The phytase activity may be determined using sodium phytate as substrate. When subjected to the action of phytase, inorganic phosphorous is liberated from sodium phytate. In a ferrous/molybdenum containing reagent, phosphorous (PO_4) forms a complex, which can be monitored spectrophotometrically at 750 nm.

One Phytase Unit (FYT) is defined as the amount of enzyme which under standard conditions (i.e. at pH 5.5, 37°C, a substrate concentration of 5.0 mM sodium phytate, and a reaction time of 30 minutes) liberates 1 μmol of phosphate per minute.

A standard operating procedure EAL-SM-0403.01 describing this analytical method in more detail is available upon request to Novo Nordisk A/S, Denmark, which publication is hereby included by reference.

Protease Activity (AU)

The proteolytic activity may be determined with denatured hemoglobin as substrate. In the Anson-Hemoglobin method for the determination of proteolytic activity denatured hemoglobin is digested, and the undigested hemoglobin is precipitated with trichloroacetic acid (TCA). The amount of TCA soluble

product is determined with phenol reagent, which gives a blue color with tyrosine and tryptophan.

One Anson Unit (AU) is defined as the amount of enzyme which under standard conditions (i.e. 25°C, pH 7.5 and 10 min. reaction time) digests hemoglobin at an initial rate such that there is liberated per minute an amount of TCA soluble product which gives the same color with phenol reagent as one milliequivalent of tyrosine.

A folder AF 4/5 describing the analytical method in more detail is available upon request to Novo Nordisk A/S, Denmark, which folder is hereby included by reference.

EXAMPLES

The invention is further illustrated with reference to the following examples which are not intended to be in any way limiting to the scope of the invention as claimed.

15 EXAMPLE 1

Soy Protein Hydrolysis

Soy protein concentrate (Unico 75, Loders Crooklaan, NL) was suspended in deionized water at 50°C.

The mixture was heat treated at 85°C for 3 minutes and cooled to 55°C, the temperature of hydrolysis. pH was adjusted to 8,5 using 4N NaOH.

The mixture was divided into three vessels for three comparative experiments:

- 1) Hydrolysing with:
Alcalase™ 2.4L, dosage 2 w/w% of protein content.
Neutrase™ 0.5L, dosage 1 w/w% of protein content.

Phytase Novo™ (an *Aspergillus* phytase, 6900 FYT/g), dosage 1 mg/g soy concentrate. pH adjusted to 4.5 with HCl. Phytase is added when pH < 6,6.

2)

Hydrolysing with:

5

Alcalase™ 2.4L, dosage 2 w/w% of protein content.

Neutrase™ 0.5L, dosage 1 w/w% of protein content.

3)

pH was adjusted to 6,2 using HCl. Phytase Novo™ (an *Aspergillus* phytase, 6900 FYT/g), dosage 1 mg/g soy concentrate.

The three hydrolyses were followed by monitoring the degree of hydrolysis and the osmolality. The results are presented in Tables 1-2, below.

Table 1

Osmolality, mOsm/kg.

Time (minutes)	Exp. 1)	Exp. 2)	Exp. 3)
0	103	102	103
15	200	173	145
35	211	183	147
65	232	201	139
125	241	208	142
185	250	221	140

Table 2

Degree of Hydrolysis, % DH

Time (minutes)	Exp. 1)	Exp. 2)	Exp. 3)
20	12.2	11.9	-1.9
40	13.9	12.6	-1.6
70	13.2	12.8	-2.0
130	17.3	15.0	-2.6
190	19.5	18.2	-2.4

Both measurements of the degree of hydrolysis (%DH) show an increased degradation of the substrate. Experiment 1 yielded more solubilized protein (78.8%) when compared to Experiment 2 (77.3%).

EXAMPLE 2**In Vitro N-digestibility of Soy Bean Meal**

Two samples were prepared:

Sample A:

1 g of soybean meal

Sample B:

1 g of soybean meal with addition of 0.032 g Phytase Novo™
(an *Aspergillus* phytase, 5000 FYT/g)

Each sample was incubated with pepsin at pH 2.4 for 2 hours followed by incubation with pancreatin at pH 6.8 for 16 hours. Suspended but not digested protein was precipitated with sulfosalicylic acid. The undigested protein (as well as other undigested components) was collected by filtration and drying. The dry

matter of the filter cake and of the sample was determined, and the amount of nitrogen (N) was determined by the Kjeldahl method.

The N digestibility was calculated as follows:

$$\frac{N_{\text{sample}} \cdot \text{Dry matter}_{\text{sample}} - N_{\text{filtercake}} \cdot \text{dry matter}_{\text{filtercake}}}{N_{\text{sample}} \cdot \text{Dry matter}_{\text{sample}}} \cdot 100$$

The protein digestibility was calculated by multiplying the nitrogen (N) digestibility by 6.25.

The following protein digestibility results were obtained:

Sample A (without phytase):88,4%

Sample B (with phytase):89,9%

EXAMPLE 3

Effect of Phytase on Apparent Nitrogen Digestibility and Nitrogen Utilization in Pigs

Balance trials on pigs (weight 46-52 kg) on a barley-
wheat-soy diet.

Diet composition:

Barley 50.8%

Wheat 20.0%

Soy 24.0%

Animal fat 2.0%

Melasse 1.0%

Minerals, vitamins

and amino acids 2.2%

No addition of inorganic phosphate to the diet. The diet was pelletized at a temperature of above 60°C.

Balance trial was conducted with an adaption period of 5 days and a 7 days collection period.

12 pigs were divided into two groups. The first group was fed the diet and the second group was fed the same diet but with addition of 20.3 g/100 kg feed of Phytase Novo™ (an *Aspergillus* phytase, 7370 FYT/g).

The content of nitrogen in the diet, faeces and urine was measured and the apparent nitrogen digestibility and nitrogen utilization was calculated.

The results are presented in Table 3, below.

Table 3

Nitrogen Digestibility and Nitrogen Utilization

Nitrogen (g/day)	Diet without Phytase	Diet with Phytase	S.E.M.
Consumed	44.5	44.5	
Faeces	6.8	6.2	0.1*
Urine	20.3	19.5	0.8 NS
Deposit	17.4	18.7	0.6 NS
Digested	37.6	38.3	0.1*
Digested, %	84.7	86.1	0.3*

* $p \leq 0.01$

NS (not significant): $p > 0.05$

S.E.M.: standard error mean

CLAIMS

1. A method for the solubilization of proteins in a vegetable protein source, which method comprises the steps of
 - (a) treating the vegetable protein source with an efficient amount of one or more phytase enzymes; and
 - (b) treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes.
2. The method according to claim 1, wherein the steps of (a) and (b) are carried out simultaneous.
3. The method according to either of claims 1-2, which method further comprises the step of
 - (c) treating the vegetable protein source with an efficient amount of one or more lipolytic enzymes (EC 3.1.1.3) and/or one or more glycosidase enzymes (EC 3.2).
4. The method according to claim 3, in which the glucosidase enzyme is an amylase, in particular an α -amylase or a β -amylase, a cellulase, in particular an endo-1,4- β -glucanase (EC 3.2.1.4) or an endo-1,3- β -glucanase (3.2.1.6), a xylanase, in particular an endo-1,4- β -glucanase (EC 3.2.1.8) or a xylan-endo-1,3- β -xylosidase (EC 3.2.1.32), an α -galactosidase (EC 3.2.1.22), a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), an endoglucanase, in particular an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).
5. The method according to claim 3, in which the glucosidase enzyme is a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), or an endoglucanase, preferably an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-

1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

6. The method according to any of claims 3-5, wherein the steps of (a), (b) and (c) are carried out simultaneous.

5 7. The method according to any of claims 1-6, wherein the phytase of step (a) is a vegetable phytase, a fungal phytase, a bacterial phytase or a phytase producible by a yeast.

8. The method according to claim 7, wherein the fungal phytase is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus niger*, a strain
10 of *Aspergillus ficuum*, a strain of *Aspergillus awamori*, a strain of *Aspergillus oryzae*, a strain of *Aspergillus terreus*, or a strain of *Aspergillus nidulans*.

9. The method according to claim 7, wherein the bacterial phytase is derived from a strain of *Bacillus* or *Pseudomonas*, preferably a strain of *Bacillus subtilis*.

15 10. The method according to claim 7, wherein the phytase is derived from a yeast of the genera *Kluyveromyces* or *Saccharomyces*, preferably a strain of *Saccharomyces cerevisiae*.

11. The method according to any of claims 1-10, wherein the vegetable protein source is treated with a phytase of step (a) in an amount of from about 2 to
20 about 50000 Phytase Units (FYT) per kg of protein source, preferably of from about 50 to about 30000 FYT per kg of protein source, most preferred of from about 100 to about 10000 FYT per kg of protein source.

12. The method according to any of claims 1-11, wherein the proteolytic enzyme of step (b) is a microbial protease, a fungal protease, or pepsin or trypsin.

13. The method according to claim 12, wherein the microbial protease is derived from a strain of *Bacillus*, preferably a strain of *Bacillus licheniformis*, and/or a strain of *Bacillus subtilis*.
14. The method according to claim 12, wherein the fungal protease is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus aculeatus*, a strain of *Aspergillus niger*, a strain of *Aspergillus oryzae*.
15. The method according to any of claims 1-14, wherein the vegetable protein source is treated with one or more proteolytic enzymes of step (b) in an amount of from about 0.0001 to about 0.5 AU per kg of protein source, preferably of from about 0.001 to about 0.05 AU per kg of protein source, more preferred of from about 0.005 to about 0.03 AU per kg of protein source.
16. The method according to any of claims 1-15, wherein the vegetable protein source is a legume, a composite plant, a crucifera, or a cereal, or mixtures thereof.
17. The method according to any of the claims 1-15, wherein the vegetable protein source is soy bean.
18. The method according to any of claims 1-17, wherein the enzymatic treatment is carried out at a pH of from 3 to 9, preferably a pH of from 4 to 7.
19. The method according to any of claims 1-18, wherein the enzymatic treatment is carried out at a temperature of from about 10 to about 70°C, preferably a temperature of from about 35 to about 65°C.
20. The method according to any of claims 1-19, which method comprises the steps of
- i) suspending a vegetable protein source in water;

- ii) enzymatic treatment of the aqueous suspension of step i) with a phytase and a proteolytic enzyme;
- iii) inactivating the enzymes of step ii); and optionally
- iv) separating the solubilized protein from the aqueous suspension.

5 21. An animal feed additive comprising a phytase and a proteolytic enzyme.

22. The animal feed additive according to claim 21, which comprises one or more additional enzymes selected from the group consisting of lipolytic enzymes (EC 3.1.1.3) and glycosidase enzymes (EC 3.2).

10 23. The animal feed additive according to claim 22, in which the glucosidase enzyme is an amylase, in particular an α -amylase or a β -amylase, a cellulase, in particular an endo-1,4- β -glucanase (EC 3.2.1.4) or an endo-1,3- β -glucanase (3.2.1.6), a xylanase, in particular an endo-1,4- β -glucanase (EC 3.2.1.8) or a xylan-endo-1,3- β -xylosidase (EC 3.2.1.32), an α -galactosidase (EC 3.2.1.22),
15 a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), an endoglucanase, in particular an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

24. The animal feed additive according to claim 22, in which the
20 glucosidase enzyme is a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), or an endoglucanase, preferably an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

25 25. The animal feed additive according to any of claims 21-24, wherein the phytase is a vegetable phytase, a fungal phytase, a bacterial phytase, or a phytase producible by a yeast.

26. The animal feed additive according to claim 25, wherein the fungal phytase is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus niger*, a strain of *Aspergillus ficuum*, a strain of *Aspergillus awamori*, a strain of *Aspergillus oryzae*, a strain of *Aspergillus terreus*, or a strain of *Aspergillus nidulans*.

27. The animal feed additive according to claim 25, wherein the bacterial phytase is derived from a strain of *Bacillus* or a strain of *Pseudomonas*, preferably a strain of *Bacillus subtilis*.

28. The animal feed additive according to claim 25, wherein the phytase is derived from a yeast of the genus *Kluyveromyces* or the genus *Saccharomyces*, preferably a strain of *Saccharomyces cerevisiae*.

29. The animal feed additive according to any of claims 21-28, comprising one or more phytase enzymes in an amount corresponding to of from about 200 to about 50000 Phytase Units (FYT) per gram of the composition, preferably of from about 500 to about 10000 FYT per gram of the composition, more preferred of from about 2000 to about 6000 FYT per gram of the composition.

30. The animal feed additive according to any of claims 21-29, wherein the proteolytic enzyme is a microbial protease, a fungal protease, or pepsin or trypsin.

31. The animal feed additive according to claim 30, wherein the microbial protease is derived from a strain of *Bacillus*, preferably a strain of *Bacillus licheniformis* or a strain of *Bacillus subtilis*.

32. The animal feed additive according to claim 30, wherein the fungal protease is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus aculeatus*, a strain of *Aspergillus niger*, or a strain of *Aspergillus oryzae*.

33. The animal feed additive according to any of claims 21-32, comprising a proteolytic enzyme in an amount corresponding to of from about 0.01 to about 20 AU (Anson Units) per gram of the composition, preferably of from about 0.1 to about 10 AU per gram of the composition, more preferred of from about 0.2 to about 5 AU per gram of the composition.

34. The animal feed additive according to any of the claims 21-33, wherein one or more of the enzymes are provided in the form of a powder, a non-dusting granulate, a stabilized liquid, a slurry, or as protected enzymes.

AMENDED CLAIMS

[received by the International Bureau on 21 September 1995 (21.09.95);
original claims 2 and 20 cancelled; claims 1 and 3 replaced by
amended claim 1; claims 4-19, 21 and 23-34 replaced by new claims
2-17, 18 and 19-30 (5 pages)]

1. A method for the solubilization of proteins in a vegetable protein source, which method comprises the steps of
 - (a) treating the vegetable protein source with an efficient amount of one or more phytase enzymes;
 - (b) treating the vegetable protein source with an efficient amount of one or more proteolytic enzymes; and
 - (c) treating the vegetable protein source with an efficient amount of one or more lipolytic enzymes (EC 3.1.1.3) and/or one or more glycosidase enzymes (EC 3.2).
2. The method according to claim 1, in which the glucosidase enzyme of step (c) is an amylase, in particular an α -amylase or a β -amylase, a cellulase, in particular an endo-1,4- β -glucanase (EC 3.2.1.4) or an endo-1,3- β -glucanase (3.2.1.6), a xylanase, in particular an endo-1,4- β -glucanase (EC 3.2.1.8) or a xylan-endo-1,3- β -xylosidase (EC 3.2.1.32), an α -galactosidase (EC 3.2.1.22), a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), an endoglucanase, in particular an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).
3. The method according to claim 1, in which the glucosidase enzyme of step (c) is a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), or an endoglucanase, preferably an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).
4. The method according to any of claims 1-3, wherein the steps of (a), (b) and (c) are carried out simultaneous.

5. The method according to any of claims 1-4, wherein the phytase of step (a) is a vegetable phytase, a fungal phytase, a bacterial phytase or a phytase producible by a yeast.
6. The method according to claim 5, wherein the fungal phytase is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus niger*, a strain of *Aspergillus ficuum*, a strain of *Aspergillus awamori*, a strain of *Aspergillus oryzae*, a strain of *Aspergillus terreus*, or a strain of *Aspergillus nidulans*.
7. The method according to claim 5, wherein the bacterial phytase is derived from a strain of *Bacillus* or *Pseudomonas*, preferably a strain of *Bacillus subtilis*.
8. The method according to claim 5, wherein the phytase is derived from a yeast of the genera *Kluyveromyces* or *Saccharomyces*, preferably a strain of *Saccharomyces cerevisiae*.
9. The method according to any of claims 1-8, wherein the vegetable protein source is treated with a phytase of step (a) in an amount of from about 2 to about 50000 Phytase Units (FYT) per kg of protein source, preferably of from about 50 to about 30000 FYT per kg of protein source, most preferred of from about 100 to about 10000 FYT per kg of protein source.
10. The method according to any of claims 1-9, wherein the proteolytic enzyme of step (b) is a microbial protease, a fungal protease, or pepsin or trypsin.
11. The method according to claim 10, wherein the microbial protease is derived from a strain of *Bacillus*, preferably a strain of *Bacillus licheniformis*, and/or a strain of *Bacillus subtilis*.

12. The method according to claim 10, wherein the fungal protease is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus aculeatus*, a strain of *Aspergillus niger*, a strain of *Aspergillus oryzae*.
13. The method according to any of claims 1-12, wherein the vegetable protein source is treated with one or more proteolytic enzymes of step (b) in an amount of from about 0.0001 to about 0.5 AU per kg of protein source, preferably of from about 0.001 to about 0.05 AU per kg of protein source, more preferred of from about 0.005 to about 0.03 AU per kg of protein source.
14. The method according to any of claims 1-13, wherein the vegetable protein source is a legume, a composite plant, a crucifera, or a cereal, or mixtures thereof.
15. The method according to any of the claims 1-13, wherein the vegetable protein source is soy bean.
16. The method according to any of claims 1-15, wherein the enzymatic treatment is carried out at a pH of from 3 to 9, preferably a pH of from 4 to 7.
17. The method according to any of claims 1-16, wherein the enzymatic treatment is carried out at a temperature of from about 10 to about 70°C, preferably a temperature of from about 35 to about 65°C.
18. An animal feed additive comprising a phytase, a proteolytic enzyme, and one or more additional enzymes selected from the group consisting of lipolytic enzymes (EC 3.1.1.3) and glycosidase enzymes (EC 3.2).
19. The animal feed additive according to claim 18, in which the glucosidase enzyme is an amylase, in particular an α -amylase or a β -amylase, a cellulase, in particular an endo-1,4- β -glucanase (EC 3.2.1.4) or an endo-1,3- β -

glucanase (3.2.1.6), a xylanase, in particular an endo-1,4- β -glucanase (EC 3.2.1.8) or a xylan-endo-1,3- β -xylosidase (EC 3.2.1.32), an α -galactosidase (EC 3.2.1.22), a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), an endoglucanase, in particular an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

20. The animal feed additive according to claim 18, in which the glucosidase enzyme is a polygalacturonase (EC 3.2.1.15), a cellulose-1,4- β -cellobiosidase (EC 3.2.1.91), or an endoglucanase, preferably an endo-1,6- β -glucanase (EC 3.2.1.75), an endo-1,2- β -glucanase (EC 3.2.1.71), an endo-1,3- β -glucanase (EC 3.2.1.39) or an endo-1,3- α -glucanase (EC 3.2.1.59).

21. The animal feed additive according to any of claims 18-20, wherein the phytase is a vegetable phytase, a fungal phytase, a bacterial phytase, or a phytase producible by a yeast.

15 22. The animal feed additive according to claim 21, wherein the fungal phytase is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus niger*, a strain of *Aspergillus ficuum*, a strain of *Aspergillus awamori*, a strain of *Aspergillus oryzae*, a strain of *Aspergillus terreus*, or a strain of *Aspergillus nidulans*.

23. The animal feed additive according to claim 21, wherein the bacterial 20 phytase is derived from a strain of *Bacillus* or a strain of *Pseudomonas*, preferably a strain of *Bacillus subtilis*.

24. The animal feed additive according to claim 21, wherein the phytase is derived from a yeast of the genus *Kluyveromyces* or the genus *Saccharomyces*, preferably a strain of *Saccharomyces cerevisiae*.

25. The animal feed additive according to any of claims 18-24, comprising one or more phytase enzymes in an amount corresponding to of from about 200 to about 50000 Phytase Units (FYT) per gram of the composition, preferably of from about 500 to about 10000 FYT per gram of the composition, more preferred of from s about 2000 to about 6000 FYT per gram of the composition.
26. The animal feed additive according to any of claims 18-25, wherein the proteolytic enzyme is a microbial protease, a fungal protease, or pepsin or trypsin.
27. The animal feed additive according to claim 26, wherein the microbial protease is derived from a strain of *Bacillus*, preferably a strain of *Bacillus licheniformis* or a strain of *Bacillus subtilis*.
28. The animal feed additive according to claim 26, wherein the fungal protease is derived from a strain of *Aspergillus*, preferably a strain of *Aspergillus aculeatus*, a strain of *Aspergillus niger*, or a strain of *Aspergillus oryzae*.
29. The animal feed additive according to any of claims 18-28, comprising 15 a proteolytic enzyme in an amount corresponding to of from about 0.01 to about 20 AU (Anson Units) per gram of the composition, preferably of from about 0.1 to about 10 AU per gram of the composition, more preferred of from about 0.2 to about 5 AU per gram of the composition.
30. The animal feed additive according to any of the claims 18-29, wherein 20 one or more of the enzymes are provided in the form of a powder, a non-dusting granulate, a stabilized liquid, a slurry, or as protected enzymes.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 95/00166

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: A23J 3/34, A23K 1/14, A23L 1/105

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: A23J, A23K, A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3966971 A (ALPHA L. MOREHOUSE ET AL), 29 June 1976 (29.06.76), column 2, line 48 - line 64, abstract --	1-34
A	WO 9207474 A1 (E.I. DU PONT DE NEMOURS AND COMPANY), 14 May 1992 (14.05.92), page 4, line 3 - line 21; page 6, line 27 - page 7, line 10 -- -----	1-34

☐ Further documents are listed in the continuation of Box C.
 ☒ See patent family annex.

- * Special categories of cited documents:
- "A" document defining the general state of the art which is not considered to be of particular relevance
- "B" earlier document but published on or after the international filing date
- "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed
- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search 24 July 1995	Date of mailing of the international search report 25 -07- 1995
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86	Authorized officer Jack Hedlund Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT
Information on patent family members

29/05/95

International application No.

PCT/DK 95/00166

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A- 3966971	29/06/76	AT-B- 352510	25/09/79
		AU-A- 8786675	30/06/77
		BE-A, A- 837267	16/04/76
		DE-A- 2600060	08/07/76
		FR-A, B- 2296373	30/07/76
		GB-A- 1507880	19/04/78
		JP-A- 51125300	01/11/76
		NL-A- 7515245	06/07/76
		SE-A- 7514773	05/07/76
WO-A1- 9207474	14/05/92	AU-A- 9016291	26/05/92